

Literatur

- [1] Weerheijm KL, Jalevik B, Alaluusua S: Molar-incisor hypomineralisation. *Caries Res* 35 (5), 390–391 (2001).
- [2] Weerheijm KL, Duggal M, Mejare I, Papagiannoulis L, Koch G, Martens LC et al.: Judgement criteria for molar incisor hypomineralisation (MIH) in epidemiologic studies: a summary of the European meeting on MIH held in Athens, 2003. *Eur J Paediatr Dent* 4 (3), 110–113 (2003).
- [3] Koch G, Hallonsten AL, Ludvigsson N, Hansson BO, Holst A, Ullbro C: Epidemiologic study of idiopathic enamel hypomineralization in permanent teeth of Swedish children. *Community Dent Oral Epidemiol* 15 (5), 279–285 (1987).
- [4] Fuchs C, Buske G, Krämer N: Schmelzbildungsstörungen – Fallbericht einer generalisierten Schmelzbildungsstörung in der 1. Dentition (Enamel malformations – Case report of a generalised enamel malformation in the primary dentition). *Oral Prophyl* 31 (4), 178–186 (2009).
- [5] Elfrink ME, ten Cate JM, Jaddoe VW, Hofman A, Moll HA, Veerkamp JS: Deciduous molar hypomineralization and molar incisor hypomineralization. *J Dent Res* 91 (6), 551–555 (2012).
- [6] Zhao D, Dong B, Yu D, Ren Q, Sun Y: The prevalence of molar incisor hypomineralization: evidence from 70 studies. *Int J Paediatr Dent* (2017).
- [7] Jalevik B, Noren JG, Klingberg G, Barregard L: Etiologic factors influencing the prevalence of demarcated opacities in permanent first molars in a group of Swedish children. *Eur J Oral Sci* 109 (4), 230–234 (2001).
- [8] Dietrich G, Sperling S, Hetzer G: Molar incisor hypomineralisation in a group of children and adolescents living in Dresden (Germany). *Eur J Paediatr Dent* 4 (3), 133–137 (2003).
- [9] Petrou MA, Giraki M, Bissar AR, Basner R, Wempe C, Altarabulsi MB et al.: Prevalence of molar-incisor-hypomineralisation among school children in four German cities. *Int J Paediatr Dent* 24 (6), 434–440 (2014).
- [10] Jordan R, Micheelis W: Fünfte Deutsche Mundgesundheitsstudie (DMS V). Kassenzahnärztliche Bundesvereinigung; Bundeszahnärztekammer (2016).
- [11] Alaluusua S: Aetiology of molar-incisor hypomineralisation: a systematic review. *Eur Arch Paediatr Dent* 11 (2), 53–58 (2010).

- [12] Crombie F, Manton D, Kilpatrick N: Aetiology of molar-incisor hypomineralization: a critical review. *Int J Paediatr Dent* 19 (2), 73–83 (2009).
- [13] Taylor GD: Molar incisor hypomineralisation. *Evid Based Dent* 18 (1), 15–16 (2017).
- [14] Beentjes VE, Weerheijm KL, Groen HJ: Factors involved in the aetiology of molar-incisor hypomineralisation (MIH). *Eur J Paediatr Dent* 3 (1), 9–13 (2002).
- [15] Silva MJ, Scurrah KJ, Craig JM, Manton DJ, Kilpatrick N: Etiology of molar incisor hypomineralization – a systematic review. *Community Dent Oral Epidemiol* 44 (4), 342–353 (2016).
- [16] Jedeon K, De la Dure-Molla M, Brookes SJ, Liodice S, Marciano C, Kirkham J et al.: Enamel defects reflect perinatal exposure to bisphenol A. *Am J Pathol* 183 (1), 108–118 (2013).
- [17] Souza JF, Costa-Silva CM, Jeremias F, Santos-Pinto L, Zuanon AC, Cordeiro RC: Molar incisor hypomineralisation: possible aetiological factors in children from urban and rural areas. *Eur Arch Paediatr Dent* 13 (4), 164–170 (2012).
- [18] Lygidakis NA, Wong F, Jalevik B, Vierrou AM, Alaluusua S, Espelid I: Best Clinical Practice Guidance for clinicians dealing with children presenting with molar-incisor-hypomineralisation (MIH): An EAPD Policy Document. *Eur Arch Paediatr Dent* 11 (2), 75–81 (2010).
- [19] Wetzel WE, Reckel U: Fehlstrukturierte Sechsjahrmolaren nehmen zu – eine Umfrage. *Zahnärztl Mitt* 81, 650–651 (1991).
- [20] Bekes K, Steffen R: Das Würzburger MIH-Konzept: Teil 1. Der MIH-Treatment Need Index (MIH-TNI). Ein neuer Index zur Befunderhebung und Therapieplanung bei Patienten mit Molaren-Inzisiven-Hypomineralisation (MIH). *Oralprophylaxe & Kinderzahnheilkunde* 38 (4), 165–170 (2016).
- [21] Bekes K, Krämer N, van Waes H, Steffen R: Das Würzburger MIH-Konzept: Teil 2. Der Therapieplan. *Oralprophylaxe & Kinderzahnheilkunde* 38 (4), 171–175 (2016).
- [22] Fayle SA: Molar incisor hypomineralisation: restorative management. *Eur J Paediatr Dent* 4 (3), 121–126 (2003).
- [23] Lygidakis NA, Chaliasou A, Siounas G: Evaluation of composite restorations in hypomineralised permanent molars: a four year clinical study. *Eur J Paediatr Dent* 4 (3), 143–148 (2003).
- [24] Jalevik B, Klingberg GA: Dental treatment, dental fear and behaviour management problems in children with severe enamel hypomineralization of their permanent first molars. *Int J Paediatr Dent* 12 (1), 24–32 (2002).

- [25] Zagdwon AM, Toumba KJ, Curzon ME: The prevalence of developmental enamel defects in permanent molars in a group of English school children. *Eur J Paediatr Dent* 3 (2), 91–96 (2002).
- [26] Feierabend S: Behandlungskonzepte bei Strukturanomalien des Zahnschmelzes und des Dentins. *Stomatologie* 111, 341–353 (2014).
- [27] Schätzle M, Patcas R: Idealer Extraktionszeitpunkt bei ersten bleibenden Molaren. Eine Literaturübersicht. *Quintessenz* 62 (12), 1631–1635 (2011).
- [28] Williams JK, Gowans AJ: Hypomineralised first permanent molars and the orthodontist. *Eur J Paediatr Dent* 4 (3), 129–132 (2003).