

- [1] Andersson H, Zaren B, Frykholm P. Low incidence of pulmonary aspiration in children allowed intake of clear fluids until called to the operating suite. *Paediatr Anaesth.* 2015; 25 (8): 770-7.
- [2] Beck CE, Rudolph D, Mahn C, Etspuler A, Korf M, Luthke M, et al. Impact of clear fluid fasting on pulmonary aspiration in children undergoing general anesthesia: Results of the German prospective multicenter observational (NiKs) study. *Paediatr Anaesth.* 2020; 30 (8): 892-9.
- [3] Bertolizio G, Saint-Martin C, Ingelmo P. Cervical instability in patients with Trisomy 21: The eternal gamble. *Paediatr Anaesth.* 2018; 28 (10): 830-3.
- [4] Bromfalk A, Myrberg T, Wallden J, Engstrom A, Hultin M. Preoperative anxiety in preschool children: A randomized clinical trial comparing midazolam, clonidine, and dexmedetomidine. *Paediatr Anaesth.* 2021; 31 (11): 1225-33.
- [5] Brown KA, Laferriere A, Lakheeram I, Moss IR. Recurrent hypoxemia in children is associated with increased analgesic sensitivity to opiates. *Anesthesiology.* 2006; 105 (4): 665-9.
- [6] Cox RG, Nemish U, Ewen A, Crowe MJ. Evidence-based clinical update: does premedication with oral midazolam lead to improved behavioural outcomes in children? *Can J Anaesth.* 2006; 53 (12): 1213-9.
- [7] Cuper NJ, de Graaff JC, van Dijk AT, Verdaasdonk RM, van der Werff DB, Kalkman CJ. Predictive factors for difficult intravenous cannulation in pediatric patients at a tertiary pediatric hospital. *Paediatr Anaesth.* 2012; 22 (3): 223-9.
- [8] Frei FJ, Haemmerle MH, Brunner R, Kern C. Minimum alveolar concentration for halothane in children with cerebral palsy and severe mental retardation. *Anaesthesia.* 1997; 52 (11): 1056-60.
- [9] Frykholm P, Disma N, Andersson H, Beck C, Bouvet L, Cercueil E, et al. Pre-operative fasting in children: A guideline from the European Society of Anaesthesiology and Intensive Care. *Eur J Anaesthesiol.* 2022; 39 (1): 4-25.
- [10] Garcia-Marcinkiewicz AG, Kovatsis PG, Hunyady AI, Olomu PN, Zhang B, Sathyamoorthy M, et al. First-attempt success rate of video laryngoscopy in small infants (VISI): a multicentre, randomised controlled trial. *Lancet.* 2020; 396 (10266): 1905-13.
- [11] Glatz P, Sandin RH, Pedersen NL, Bonamy AK, Eriksson LI, Granath F. Association of Anesthesia and Surgery During Childhood With Long-term Academic Performance. *JAMA Pediatr.* 2017; 171 (1): e163470.

- [12] Habre W, Disma N, Virag K, Becke K, Hansen TG, Jöhr M, et al. Incidence of severe critical events in paediatric anaesthesia (APRICOT): a prospective multicentre observational study in 261 hospitals in Europe. *Lancet Respir Med*. 2017; 5 (5): 412-25.
- [13] Hata T, Todd MM. Cervical spine considerations when anesthetizing patients with Down syndrome. *Anesthesiology*. 2005; 102 (3): 680-5.
- [14] Ing C, Warner DO, Sun LS, Flick RP, Davidson AJ, Vutskits L, et al. Anesthesia and Developing Brains: Unanswered Questions and Proposed Paths Forward. *Anesthesiology*. 2022; 136 (3): 500-12.
- [15] Jöhr M. *Kinderanästhesie*. 9. Auflage. Elsevier, München, 2019.
- [16] Kraemer FW, Stricker PA, Gurnaney HG, McClung H, Meador MR, Sussman E, et al. Bradycardia during induction of anesthesia with sevoflurane in children with Down syndrome. *Anesth Analg*. 2010; 111 (5): 1259-63.
- [17] Lewanda AF, Matisoff A, Revenis M, Harahsheh A, Futterman C, Nino G, et al. Preoperative evaluation and comprehensive risk assessment for children with Down syndrome. *Paediatr Anaesth*. 2016; 26 (4): 356-62.
- [18] Lonnqvist PA, Habre W. Midazolam as premedication: is the emperor naked or just half-dressed? *Paediatr Anaesth*. 2005; 15 (4): 263-5.
- [19] Lyseng-Williamson KA. Midazolam oral solution (Ozalin): a profile of its use for procedural sedation or premedication before anaesthesia in children. *Drugs & Therapy Perspectives*. 2019; 35: 255-62.
- [20] McCann ME, de Graaff JC, Dorris L, Disma N, Withington D, Bell G, et al. Neurodevelopmental outcome at 5 years of age after general anaesthesia or awake-regional anaesthesia in infancy (GAS): an international, multicentre, randomised, controlled equivalence trial. *Lancet*. 2019; 393 (10172): 664-77.
- [21] Moorthy SS, Krishna G, Dierdorf SF. Resistance to vecuronium in patients with cerebral palsy. *Anesth Analg*. 1991; 73 (3): 275-7.
- [22] Prince J, Goertzen C, Zanjir M, Wong M, Azarpazhooh A. Airway Complications in Intubated Versus Laryngeal Mask Airway-Managed Dentistry: A Meta-Analysis. *Anesth Prog*. 2021; 68 (4): 193-205.
- [23] Pringle B, Dahlquist LM, Eskenazi A. Memory in pediatric patients undergoing conscious sedation for aversive medical procedures. *Health Psychol*. 2003; 22 (3): 263-9.
- [24] Sun LS, Li G, Miller TL, Salorio C, Byrne MW, Bellinger DC, et al. Association Between a Single General Anesthesia Exposure Before Age 36 Months and Neurocognitive Outcomes in Later Childhood. *JAMA*. 2016; 315 (21): 2312-20.
- [25] Warner DO, Zaccariello MJ, Katusic SK, Schroeder DR, Hanson AC, Schulte PJ, et al. Neuropsychological and Behavioral Outcomes after Exposure of Young Children

to Procedures Requiring General Anesthesia: The Mayo Anesthesia Safety in Kids (MASK) Study. *Anesthesiology*. 2018; 129 (1): 89-105.

- [26] Weiss M, Machotta A. Qualität und sichere Anästhesie für alle Kinder: Das ist ihr Recht! *Anaesthesist*. 2022; 71 (4): 255-63.
- [27] Weiss M, Vutskits L, Hansen TG, Engelhardt T. Safe Anesthesia For Every Tot - The SAFETOTS initiative. *Curr Opin Anaesthesiol*. 2015; 28 (3): 302-7.
- [28] Zickerman C, Hult AC, Hedlund L, Winso O, Johansson G, Haney M. Clonidine Versus Midazolam Premedication and Postoperative Negative Behavioral Changes in Younger Children: A Randomized Controlled Trial. *Anesth Analg*. 2022 Feb 24 ePub ahead of Print.