

**Gesunde Zähne – ein Leben lang
3-fach-Prophylaxe für Jugendliche**

Literatur

- [1] American Academy of Pediatric Dentistry. Adolescent Oral Health Care. Recommendations: Best Practices, 2015;221–228.
- [2] Salas MMS, Nascimento GG, Huysmans MC. Estimated prevalence of erosive tooth wear in permanent teeth of children and adolescents: an epidemiological systematic review and meta-regression analysis. J Dent. 2015;43:42–50.
- [3] Kirkham J, Robinson C, Strong M, Shore RC. Effects of frequency of acid exposure on demineralization/remineralization behavior of human enamel in vitro. Caries Res 1994;28(1):9–13.
- [4] Bartsch S: Jugend Isst Anders. UGB-Forum 5/11:214–217.
<https://www.fairberaten.net/saeuglings-und-kinderernaehrung/hintergrundinfos/jugend-isst-anders/druckansicht.pdf>
(aufgerufen am 21. Juni 2019)
- [5] Fine DH et al. Effect of rinsing with an essential oil-containing mouthrinse on subgingival periodontopathogens. J Periodontol 2007;78:1935–42.
- [6] Lux R et al. data on file. Confocal laser scanning microscopy evaluation of biofilm kill activity of representative global mouthrinse. 2005.
- [7] Sharma N et al. Adjunctive benefit of an essential oil-containing mouthrinse in reducing plaque and gingivitis in patients who brush and floss regularly: a six-month study. JADA 2004; 135: 496–504.
- [8] Fine DH et al. Effect of rinsing with an essential oil-containing mouthrinse on subgingival periodontopathogens. J Periodontol 2007;78:1935-1942.
- [9] Sharma N et al. Adjunctive benefit of an essential oil-containing mouthrinse in reducing plaque and gingivitis in patients who brush and floss regularly: a six-month study. J Am Dent Assoc 2004;135(4):496–504.